


SJRH ED COVID-19 Obstetrical Patient Care Algorithm

- * Medically stable OB patients should be managed in L&D AND Active Medical/non-OB issues in ED*
- *** Moderate/severely symptomatic patients >20wGA AND pregnancy-related emergency must be stabilized in ED then transferred to L&D with droplet-contact precautions. Attending OBGYN on-call notified STAT ***
- * Mild patients with >20wGA AND obstetrical emergency may be transferred to L&D with droplet-contact precautions. Attending OBGYN on-call and L&D notified STAT *
- ** Patients in active labour require rapid COVID swab **




Triage to COVID Cove or Negative Pressure Room appropriate to clinical presentation * see page 2

Attending provider to notify attending OBGYN on-call of any pregnant woman >20w GA in ED who screens positive for COVID.

- o L&D RN and fetal surveillance equipment can be available for ED at discretion of OBGYN

- Clinical Management Pearls
 - o Assess FHR
 - o **Target SpO2>95%**
 - o Avoid supine positioning – left lateral tilt with right hip wedge must be employed, especially if ALC
 - o Conscious prone positioning – extra caution and attention and under co-consultation with ICU/OBGYN
- Critically ill: best fetal resuscitation is optimizing maternal resuscitation. See SJRH COVID-19 Clinical Management Guidance
- Consult Internist/ICE on-call re: admission of moderate/severely symptomatic PUI COVID. Co-consult OB for >20w gestation
- Obstetrical admissions to 3AN/LD per attending OBGyn

*** if at any time, patients >20w gestation are deemed to be in active labor or have another pregnancy-related emergency (SRM, active labor, PVB, abruption, etc.) efforts should be made to stabilize in ED, then immediately transfer to labor & delivery unit under droplet-contact precautions. OBGYN on-call should be notified STAT ***



Attending provider to notify attending OBGYN on-call of any pregnant woman >20w GA in ED who screens positive for COVID.

- o L&D RN and fetal surveillance equipment can be available for ED at discretion of OBGYN
- Clinical Management Pearls
 - o Assess FHR
 - o **Target SpO₂>95%**
 - o Avoid supine positioning – left lateral tilt with right hip wedge must be employed, especially if ALC
 - o Conscious prone positioning – extra caution and attention and under co-consultation with ICU/OBGYN
- Critically ill: best fetal resuscitation is optimizing maternal resuscitation. See SJRH COVID-19 Clinical Management Guidance
- Consult Internist/ICE on-call re: admission of moderate/severely symptomatic PUI COVID. Co-consult OB for >20w gestation.
- Obstetrical admissions to 3AN/LD per attending OBGyn

***** if at any time, patients >20w gestation are deemed to be in active labor or have another pregnancy-related emergency (SRM, active labor, PVB, abruption, etc.) efforts should be made to stabilize in ED, then immediately transfer to labor & delivery unit under droplet-contact precautions. OBGYN on-call should be notified STAT *****

***** Patients in active labor require rapid COVID-19 swab *****